

## Colon Prep: GoLytey Full Extra Dose (2-Gallons)

(Follow same instructions for Nulytey/Trilyte/Gavilyte/Peg 3350 Substitutes)

\*Google "WWMG GI Pre-Procedure Instructions" to locate this instruction PDF file & to watch the instruction video.

| General Instructions  | 5 days before  | One day before   | Colonoscopy day:   |
|---|--|--|--|
| <p>* <b>Read bowel prep instruction and watch instruction video 1 week prior to colonoscopy.</b></p> <p><b>LATE CANCELLATION FEE \$250</b>, if cancelled within 5 business days.</p> <p>Secure a <b>driver</b> for the procedure day, no driving for 12 hrs. after procedure.</p> <p>Check insurance for requirement of referral or pre-approval.</p> <p><b>If you have implanted devices</b> such as pacemaker, defibrillator, sleep apnea stimulator, neurostimulator, etc. Please notify the office.</p> <p><b>If you have an electronic stimulator</b>, bring your remote control to turn off your stimulator.</p> <p><b>If you have an ostomy</b>, please bring spare ostomy supplies with you.</p> <p>Do not bring jewelry, money, and other valuables to your colonoscopy.</p> <p><b>SUPPLIES:</b></p> <ul style="list-style-type: none"> <li>• <b>2 Gallons - GoLytey or a generic equivalent (prescription)</b></li> <li>• <b>Infant Mylicon Gas Relief Drops (simethicone)</b><br/>0.5 oz (15 mL) box</li> <li>• MiraLAX (OTC), optional</li> <li>• Fleet enema (OTC), optional</li> </ul> <p><b>Questions: call (425) 259-3122</b></p> | <ul style="list-style-type: none"> <li>• Start <b>low fiber diet</b> now.</li> <li>• <b>Read the instructions now.</b></li> <li>• <b>Watch</b> bowel preparation <b>video now*</b></li> <li>• <b>Stop following medications now:</b> oil supplements, oil capsules, and iron pills.</li> </ul> <p><b>Other medication instructions:</b></p> <ul style="list-style-type: none"> <li>• If you're taking <b>diabetic medications</b>, please ask for special instructions.</li> <li>• If you're taking <b>weight loss medications</b>, please ask for special instructions.</li> <li>• If you're taking <b>blood thinners</b>, such as Plavix, Effient, Warfarin, Eliquis, Pradaxa, or Xarelto, etc. please ask for special instructions.</li> </ul> <p><b>If you have constipation:</b></p> <ul style="list-style-type: none"> <li>• Take OTC <b>MiraLAX</b> one dose in a 8 oz glass of water up to 3 times per day as needed to maintain daily soft BMs.</li> <li>• Also ask for an <b>extra dose prep</b> regimen.</li> </ul> | <p><b>AM:</b> Mix both GoLytey containers with 1 gallon of water each and store in refrigerator.</p> <p><b>Clear liquid diet only (no solid foods!) for Entire Day:</b> clear fruit juice without pulp such as apple juice, white grape juice, broth, Gatorade (No Gatorade Frost), popsicles, small amounts of coffee or tea without cream. No milk, red, blue, or purple colors.</p> <p><b>If you take diabetic medication(s):</b> follow special instructions.</p> <p><b>8 AM:</b> Drink half of the 1st gallon of GoLytey, 8 oz glass every 15 to 20 minutes as tolerated for a total of 8 glasses. Keep remaining for 12 pm dosing.</p> <p><b>12 PM:</b> Drink the remaining GoLytey from the 1st gallon, 8 oz glass every 15 to 20 minutes as tolerated for a total of 8 glasses.</p> <p><b>6 PM:</b> Drink 2nd gallon of GoLytey, 8 oz glass every 15 to 20 minutes as tolerated. Finish 10 glasses and save the other 6 glasses for the morning.</p> <p><b>What to expect:</b></p> <ul style="list-style-type: none"> <li>• It may take several hours before you have lots of diarrhea.</li> <li>• If you have nausea or vomiting, slow down the pace of drinking.</li> </ul> <p>• <b>NO ALCOHOL</b></p> | <p><b>SIX (6) HOURS PRIOR TO YOUR CHECK IN TIME:</b></p> <ul style="list-style-type: none"> <li>• Add <b>3 mL Mylicon</b> to the remaining GoLytey solution by using the syringe included in the Mylicon box. Draw up 0.6 mL per syringe and do it five times, which will give 3 mL total.</li> <li>• Drink the remaining 6 glasses of GoLytey, one 8 oz glass every 15 to 20 minutes until it is gone.</li> </ul> <p>A good prep should result in urine-like CLEAR yellowish watery stool. Complete the bowel prep even if stool looks clear halfway.</p> <p><b>Prior to 4 hours before check-in time:</b></p> <ul style="list-style-type: none"> <li>• OK to drink water or clear juice.</li> <li>• OK to take <b>usual prescribed medications</b>, unless otherwise instructed for blood thinners, diabetic &amp; weight loss medicines, etc.</li> </ul> <p><b>Within 4 hours before check-in time:</b></p> <ul style="list-style-type: none"> <li>• <b>Nothing by mouth</b>, including gum, hard candy, chewing tobacco, water, etc.</li> </ul> <p><b>Be sure to bring:</b></p> <ul style="list-style-type: none"> <li>• Medication list</li> <li>• A driver</li> <li>• Photo ID/Insurance card</li> <li>• Eyeglasses/Hearing aids</li> </ul> <p>You CANNOT drink alcohol or drive for 12 hours after the procedure.</p> |

# LOW FIBER DIET

**GENERAL RULE: DO NOT** eat anything with any of the following ingredients: nuts, seeds, raw fruits with seeds, raw vegetables, corn, beans (example: kidney, pinto, lima), whole grains, or popcorn.

## BELOW ARE THE ONLY ALLOWED FOODS

### BREADS ETC:

- White breads, rolls, biscuits
- Waffles, French toast, pancakes
- White rice, white pasta, peeled/cooked potatoes
- Saltines or white crackers, without seeds
- Hot cereal: cream of rice, cream of wheat
- Cold cereal: Rice Krispies, Corn Flakes, Special K
- 

### VEGETABLES AND FRUITS:

- Cooked and canned vegetables: carrots, green beans, spinach
- Applesauce
- Apple or white grape juice
- Bananas
- Cantaloupe, honeydew melon

### MILK/DAIRY:

- Milk: plain or flavored, coconut, almond, soy
- Yogurt, custard and ice cream
- Cheese, cottage cheese

### MEAT/PROTEIN:

- Well cooked, lean meats, fish
- Eggs
- Creamy peanut/almond butter
- Tofu

### FATS, SWEETS, CONDIMENTS, BEVERAGES:

- Margarine, butter, cooking oils, mayonnaise, salad dressing
- Plain gravies
- Sugar, clear jelly (seedless), honey and syrup
- Soups made with allowed vegetables, broth
- Coffee, tea, carbonated beverages
- Plain cake
- Gelatin, plain pudding, popsicles
- Hard candy, pretzels
- Plain chocolate