

# **Colon Prep: SUPREP**

Google "WWMG GI Pre-Procedure Instructions" to locate this instruction PDF file.

General Instructions	5 days before	One day before	Colonoscopy day:
Read bowel prep instruction 1 week prior to colonoscopy.	• Start low fiber diet now. • Read the instructions now.	DO NOT FOLLOW INSTRUCTIONS INSIDE/ON THE BOX	SIX (6) HOURS PRIOR TO YOUR CHECK IN TIME:
LATE CANCELLATION FEE \$250, if cancelled within 5 business days.	• Stop following medications now: oil supplements, oil capsules, and iron pills.	Clear liquid diet only (no solid foods!) for Entire Day: clear fruit juice without pulp such as apple juice, white grape	Follow exact 6 PM instructions from the day before. Add <b>3 mL Mylicon</b> to the above solution by using the syringe included in the
Secure a <b>driver</b> for the procedure day, no driving for 12 hrs. after procedure.	Other medication instructions: • If you're taking diabetic medications, please ask for special	juice, broth, Gatorade (No Gatorade Frost), popsicles, small amounts of coffee or tea without cream. No milk,	Mylicon box. Draw up 0.6 mL per syringe and do it five times, which will give 3 mL total. Drink the 2nd 6 oz SUPREP bottle.
Check insurance for requirement of referral or pre-approval.	instructions.  • If you're taking weight loss	red, blue, or purple colors.	A good prep should result in urine-like CLEAR yellowish watery stool. Complete the
<b>If you have implanted devices</b> such as pacemaker, defibrillator, sleep	medications, please ask for special instructions.	If you take diabetic medication(s): follow special instructions.	bowel prep even if stool looks clear halfway.
apnea stimulator, neurostimulator, etc. Please notify the office.	• If you're taking <b>blood thinners,</b> such as Plavix, Effient, Warfarin,	4 PM: Take 4 tablets of Dulcolax (Bisacodyl) Laxative 5 mg (OTC)	Prior to 4 hours before check-in time:  OK to drink water or clear juice.  OK to take years processived medications.
If you have an electronic stimulator, bring your remote control to turn off	Eliquis, Pradaxa, or Xarelto, etc. please ask for special instructions.	6 PM: Pour ONE (1) 6oz bottle of SUPREP liquid into the 16 oz container	OK to take usual prescribed medications, unless otherwise instructed for blood thinners, diabetic & weight loss medicines,
your stimulator.  If you have an ostomy, please bring	If you have constipation:  • Take OTC MiraLAX one dose in a 8	(cup). Add cold water to the 16 oz container (cup) fill line. <b>Drink 8 oz (</b> ½	etc.
spare ostomy supplies with you.	oz glass of water up to 3 times per day as needed to maintain daily soft	the cup), Repeat 15 minutes later, drink the remaining 8 oz (½ the cup)	<ul><li>Within 4 hours before check-in time:</li><li>Nothing by mouth, including gum, hard</li></ul>
Do not bring jewelry, money, and other valuables to your colonoscopy.	BMs. • Also ask for an extra dose prep	of SUPREP. You must drink (2) two more 16 oz containers (cups) of water	candy, chewing tobacco, water, etc.
<ul> <li>SUPPLIES:</li> <li>SUPREP (prescription) 1 Box</li> <li>Dulcolax Laxative (Bisacodyl) 5 mg, 4 tablets (OTC)</li> </ul>	regimen.	<ul> <li>over the next hour to stay hydrated.</li> <li>What to expect:</li> <li>It may take several hours before you have lots of diarrhea.</li> </ul>	<ul> <li>Be sure to bring:</li> <li>Medication list</li> <li>A driver</li> <li>Photo ID/Insurance card</li> <li>Eyeglasses/Hearing aids</li> </ul>
<ul> <li>Infant Mylicon Gas Relief Drops (simethicone) 0.5 oz (15mL) (OTC)</li> <li>MiraLax (OTC) optional</li> <li>Fleet enema (OTC) optional</li> <li>Questions: call (425) 259-3122</li> </ul>		<ul> <li>If you have nausea or vomiting, slow down the pace of drinking.</li> <li>NO ALCOHOL</li> </ul>	You CANNOT drink alcohol or drive for 12 hours after the procedure.



## **LOW FIBER DIET**

GENERAL RULE: DO NOT eat anything with any of the following ingredients: nuts, seeds, raw fruits with seeds, raw vegetables, corn, beans (example: kidney, pinto, lima), whole grains, or popcorn.

## BELOW ARE THE ONLY ALLOWED FOODS

#### **BREADS ETC:**

- White breads, rolls, biscuits
- Waffles, French toast, pancakes
- White rice, white pasta, peeled/cooked potatoes
- Saltines or white crackers, without seeds
- Hot cereal: cream of rice, cream of wheat
- Cold cereal: Rice Krispies, Corn Flakes, Special K

#### **VEGETABLES AND FRUITS:**

- Cooked and canned vegetables: carrots, green beans, spinach
- Applesauce
- Apple or white grape juice
- Bananas
- Cantaloupe, honeydew melon

#### MILK/DAIRY:

- Milk: plain or flavored, coconut, almond, soy
- Yogurt, custard and ice cream
- Cheese, cottage cheese

#### **MEAT/PROTEIN:**

- Well cooked, lean meats, fish
- Eggs
- Creamy peanut/almond butter
- Tofu

### **FATS, SWEETS, CONDIMENTS, BEVERAGES:**

- Margarine, butter, cooking oils, mayonnaise, salad dressing
- Plain gravies
- Sugar, clear jelly (seedless), honey and syrup
- Soups made with allowed vegetables, broth
- Coffee, tea, carbonated beverages
- Plain cake
- Gelatin, plain pudding, popsicles
- Hard candy, pretzels
- Plain chocolate